

Household and Structural Pest Management for Professionals

See PB 1303 UT Extension *Managing Pests Around the Home* (<http://www.utextension.utk.edu/publications/pests/default.asp#home>) or *The UT E&PP Redbook, Household and Structural Pests section* (<http://eppserver.ag.utk.edu/redbook/sections/structural.htm>) for information related to preventing and otherwise managing pests using integrated pest management. More information for the pest management professional can be found in the following UT Extension Publications located on the web at <http://eppserver.ag.utk.edu/PSEP/secondlevel/materials.htm>

- Vail, K.M., G. Burgess, R. Gerhardt and C. Harper [eds.]. 2007. PB1673 General Pest and Rodent Control Pesticide Applicator Licensing Manual (GRC). pp. 130.
- Vail, K., G. Burgess, R. Gerhardt and C. Jones [eds.]. 2002. PB1685 Public Health Mosquito Control: The Tennessee Mosquito Control Handbook (Pesticide Applicator Licensing Manual). pp. 54.
- Vail, K., G. Burgess, C. Pless and B. Bonds [eds.]. 2006. PB 1703 Wood-Destroying Organisms Licensing Manual.
- Vail, K.M., E. Burgess, R. Gerhardt, and Craig Harper. 2006. PB1732 Industrial, Institutional, Structural and Health Related Pest Management Certification Manual (Category 7). pp. 105
- Vail, K.M., E.E. Burgess, R. Gerhardt, C. Jones, J. Skinner and C. Harper. 2003. PB1733. Public Health Pest Control (Certification Category 8). pp. 99
- Vail, K. 2003. PB1734. Commodity and Structural Fumigation. pp. 170

Further references for household pest identification:

Manuals:

- Bennett, G.W., J.M. Owens, and R.M. Corrigan. 1997. **Truman's Scientific Guide to Pest Control Operations**. 5th. edition. Edgell Communications, Duluth, MN
- Mallis, A. 2004. **Handbook of Pest Control - the Behavior, Life History and Control of Household Pests**. 9th edition. Franzak and Foster Co. Cleveland, Ohio.

Field Guides

- Hedges, S. **Pest Control Technology Field Guide for the Management of Structure-Infesting Ants**. Franzak and Foster Co., Cleveland, Ohio.
- Hedges, S. **Pest Control Technology Field Guide for the Management of Structure-Infesting Flies**. Franzak and Foster Co., Cleveland, Ohio.
- Hedges, S. **Pest Control Technology Field Guide for the Management of Urban Spiders**. Franzak and Foster Co., Cleveland, Ohio.
- Hedges, S. and M. Lacey. **Structure-Infesting Beetles. Volume 1: Hide and Carpet beetles/Wood-Boring Beetles**. Franzak and Foster Co., Cleveland, Ohio.
- Hedges, S. and M. Lacey. **Structure-Infesting Beetles. Volume 2: Stored Product Beetles/Occasional and Overwintering Beetles**. Franzak and Foster Co.
- Klotz, J., L. Hansen, R. Pospischil and M. Rust. 2008. **Urban Ants of North America and Europe: Identification, Biology and Management**. Comstock Publishing Associates, Cornell University Press, Ithaca, New York pp. 196
- Smith, E and R. Whitman. 2007. **NPMA Field Guide to Structural Pests**. Second edition. <http://www.pestworld.org>

General identification guides for insects.

- Petersons Field Guides: Insects of North America # 19, Beetles of North America #29, Moths of Eastern North America #30

The following pages list recommended procedures and, if necessary, pesticides to manage specific pests. Remember to reduce the pest's access to food, water and shelter. If pesticides are needed, they are more likely to provide control if access to these necessities are limited. Additional publications are listed under PESTS as a PB or SP and many are now located on our web site at <http://www.utextension.utk.edu/publications/pests/default.asp#home>. Percentages that appear after the Trade Name indicate percent active ingredient. Where no percent is given, see the label for more details. SEE THE LABEL FOR ALL DIRECTIONS!

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
<p>ANTS - ARGENTINE ANT, ODOROUS HOUSE ANT</p> <p>See PB1629</p>	<p>ARGENTINE ANT Bump (Node) on waist: one, obvious Gaster tip: no circular opening Odor when crushed: "disagreeable, rotten-coconut-like" with an additional faint musty odor Color: light to dark brown</p> <p>ODOROUS HOUSE ANT Nodes on waist: one, very flat, barely noticeable node hidden by the gaster Gaster tip: no circular opening Odor when crushed: "disagreeable, rotten-coconut-like" Color: brown to black Both ant workers: 1/8-inch</p> <p>The odorous house ant is the most common house-invading ant in TN. It makes shallow nests in shady, moist areas such as mulch, pine straw, stones, logs. Moves indoors during periods of heavy rain and moves nests often. Winged male odorous house ants may be collected from lights from May through June.</p> <p>Argentine ant is not as common as the odorous house ant, but can build up larger populations. Behavior is similar for both species.</p>	<p><u>Slower-acting, Non-repellent Outdoor Perimeter Spray:</u> Terimidor SC 0.06%</p> <p><u>Outdoor Baiting:</u> Advion Mole Cricket Bait (perimeter broadcast) Gourmet Ant Bait Liquid Maxforce Fine Granule Insect Bait Maxforce Ant Killer Bait Gel (for Argentine ant) Niban FG, G baits Optigard Ant Gel Bait PRO-JOE-S Ant Bait/Gel Formula 4.5 PT 381B Advance Liquid Ant Bait PT 388B Advance Ant Gel Bait Terro- PCO (1%)</p> <p><u>Indoor Crack and Crevice Spray and Outdoor Entry Points</u> Phantom</p>	<p>Our best research results for odorous house ants have been a 0.06% fipronil (Terimidor SC) spray 1 foot up and 1 foot out from the foundation base and either COMBINED with (1) a liquid or other bait placed in the landscape near/around the structure where ants are active OR (2) Phantom applied indoors as crack and crevice in areas of activity as well as at potential entry points. Gel baits may work longer outdoors when placed in stations. Baits listed under odorous house ant and outdoor baiting have eliminated small, laboratory-maintained odorous house ant colonies within 8 weeks of bait placement or have proven effective in field studies.</p>

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
<p>ANTS - CARPENTER See PB1599</p>	<p>Large, black, red, or red-and-black ants that nest in damp wood. Wingless workers 1/4 to 3/8 inch long. Winged male and female reproductives will fly from a colony. Carpenter ants do not eat wood, but excavate smooth galleries in the wood to raise their young. Piles of coarse sawdust or splintered wood often mixed with insect parts may indicate a carpenter ant nest nearby.</p>	<p><u>Slower-acting, nonrepellent outdoor perimeter spray:</u> Termidor SC</p> <p><u>Baits:</u> <u>Indoors in cracks and crevices where ants are seen:</u> DuPont Advion Ant Bait Arena DuPont Advion Ant Gel Maxforce Carpenter Ant Bait Gel Niban FG, G</p> <p><u>Outdoors:</u> DuPont Advion Ant Bait Arena DuPont Advion Ant Gel Gourmet Ant Bait Liquid Maxforce Carpenter Ant Bait Gel Maxforce Granular Bait Niban FG, G Optigard Ant Gel Bait PT 381B Advance Liquid Ant Bait Terro -PCO (1%)</p> <p><u>Dusts:</u> Deltadust PT Perma-dust Tempo 1D</p> <p><u>Sprays (or foam according to label):</u> TalstarOne, Tempo SC Ultra, Tempo 20 WP, Cy-Kick, Transport, PT221L, Demon TC, Suspend SC, Bora-Care, Shell-guard, Tim-bor (dust, spray or foam), Armor-Guard (dust, spray or foam)</p>	<ol style="list-style-type: none"> 1. Correct moisture problem, repair leaks and ventilate. 2. If ants entering from or foraging outdoors, spray a slower acting, nonrepellent insecticide around the perimeter with an emphasis on areas where ants are entering the structure. 3. Place baits where ants are actively foraging. If baiting, do not spray ants, trails or baits with a fast-acting, repellent spray because it will kill ants too quickly and will stop them from feeding on the baits. 4. Find and treat (dust, spray or foam) nests in wood parts. <p>Drilling into the wood may be necessary. Dust nests in wall voids. Do not apply sprays near electricity.</p>
<p>ANTS-PHARAOH also called "sugar ants" or "piss ants" See PB 1629</p>	<p>Nodes on waist: two Gaster tip: sting Worker size: 1/16 - inch Antennae: 12-segmented antennae with a three-segmented club Color: yellow or orange with the end of the abdomen darkened</p> <p>Nests rarely found outdoors; however, almost any indoor crack and crevice close to sources of warmth and water.</p> <p>These ants do not swarm. Colonies multiply by "budding," in which part of an existing colony migrates, carrying brood with or without a queen to a new nesting site. Hundreds of queens and 10,000 - 100,000s workers may be present.</p>	<p><u>Baits:</u> Advion Ant Bait Arenas Maxforce Ant Killer Bait Stations Maxforce Fine Granule Insect Bait Maxforce FC Ant Bait Station Niban FG</p> <p><u>Indoor crack and crevice spray and entry points:</u> Phantom SC</p> <p><u>Outdoor entry points and trails around structure</u> Termidor SC</p>	<p>Because Pharaoh ant colonies are hidden and can occur in virtually any crack or crevice, baiting is the best way to get an insecticide back to the colony. Give a taste test of baits. Prebait entire structure with honey and natural peanut butter. Place a bait wherever ants are found. All queen(s) and all immatures must be killed to eliminate a colony. Spraying fast-acting insecticides for Pharaoh ants indoors often worsens the problem by causing the colony to split into many smaller colonies. Apply slower -acting sprays (Phantom, Termidor) around entry points during warm weather when ants are foraging outdoors.</p>

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
<p>ANTS-FIRE (see Redbook recommendations for Home Lawn Insects). Also see web sites http://www.extension.org/fire+ants for publications (Managing Fire Ants in Urban Areas, Two-step...) and other information about fire ants and "Imported Fire Ants In Tennessee" at http://fireants.utk.edu for extensive list of TN fire ant management products under "Updates". For a list of "homeowner" products by price and application type (broadcast vs. individual mound treatment) see http://www.aces.edu/pubs/docs/A/ANR-0175-A/ANR-0175-A.pdf.</p>			
<p>ANTS - OTHER See PB1629</p>	<p>Ants have elbowed antennae, a thread-like waist with one or two bumps. Unmated reproductives have wings, the front wings are larger than the hind wings. Workers ants are wingless. Ants are social insects. Colonies are usually established by a queen. Workers feed the queen, care for the brood and defend the nest. Workers travel along well-marked trails between the nest and food.</p>	<p><u>Baits:</u> DuPont Advion Ant Bait Arena DuPont Advion Ant Gel Gourmet Ant Bait Gel Gourmet Ant Bait Liquid Maxforce Ant Killer Bait Stations Maxforce Fine Granule Insect Bait Niban FG Optigard Ant Gel Bait PRO-JOE-S Ant Bait/Gel Formula 4.5 PT Advance Granular Ant Bait PT Advance 375A Select Granular PT 388B Advance Ant Gel Bait Terro-PCO (1%)</p> <p><u>Liquid Baiting Systems:</u> Gourmet Ant Bait Liquid PT 381B Advance Liquid Ant Bait Terro -PCO(1%)</p> <p><u>Pyrethroid Sprays:</u> TalstarOne, Tempo SC Ultra Tempo 20 WP Transport Suspend SC Demand CS</p> <p><u>Slower-acting, non-repellent spray:</u> Termidor SC 0.06% (outdoor perimeter) Phantom SC indoor crack and crevice, outdoor entry points</p>	<p>Follow good sanitary and exclusion practices. Exploit worker caste by using baits. Find foragers and place bait near foraging trail. Workers then bring the poisoned bait back to the nest where it is distributed among all members of the colony.</p> <p>If ants are foraging indoors from an outdoor nest, exclude ants by sealing entry points such as window sills and door steps, or spray entry points into the structure.</p>
<p>BATS See PB1624</p>	<p>Night-flying creatures invade attics and similar areas.</p>	<p>Treat area with insecticides to control external parasites including fleas, mites, and bed bugs after bats excluded.</p>	<p>Close entrance holes with wire screening to exclude bats after they have left the resting area. Seal opening with expandable foam. See details in PB 1624.</p>

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
<p>BED BUGS See PB 1763</p>	<p>Flat, oval, reddish, wingless insects. Bloodsucking. Night feeders.</p>	<p><u>Detection</u> canine detection (check for National Entomology Scent Detection Canine Association (NESCDCA) accreditation before hiring) http://www.nesdca.com/index.html</p> <p>CLIMBUP Insect interceptor, http://www.insect-interceptor.com/ (Place bed or furniture leg into center well to aid detection of new infestations and protect against reinfestation. Important to prevent other bed bug access to bed and refill wells with talc as needed.) Others</p> <p><u>Mattress and Boxspring encasements</u> Protect-A-Bed (BugLock™ 3-sided zipper system and ALLERZIP™ seal), Mattress Safes® "Ultimate" zippered encasement (zipper must be kept closed) Others</p> <p><u>Mattress</u> <i>Longer residual</i> CB-80 Suspend SC D-Force HPX</p> <p><i>Short er residual</i> Steri-Fab Bedlam PT 565 Plus XLO</p> <p><u>Combination</u> Suspend SC tank mixed with Kicker</p> <p><u>Other cracks and crevices</u> D-Force HPX Suspend SC TalstarOne Demand CS Suspend SC tank mix ed with Kicker</p> <p><u>Dusts</u> DeltaDust Drione PT Tri-Die Pressurized Silica + Pyrethrin diatomaceous earth for crawling insects Tempo 1D (remove after 4 to 6 hours)</p> <p><u>Chemical Fumigation</u> Vikane</p> <p><u>Heat fumigation</u> for potential sources of equipment see http://eppserver.ag.utk.edu/personnel/Vail/BEDBUG%20WEB%20SITES.pdf also, small chambers are available for heat fumigation</p>	<p>Treat every crack and crevices in bedroom and elsewhere, if needed. Do not spray sheets or blankets. Some PMPs will not treat mattresses with insecticides because of potential human exposure while others will cover treated mattresses and boxsprings with a plastic allergen-prevention cover after treatment. Others may physically remove bed bugs with vacuums or steam cleaners and then cover with bed encasement specifically designed to prevent bed bugs from biting through or escaping (i.e., Protect-A-Bed with BugLock™ 3-sided zipper system and ALLERZIP™ seal, or Mattress Safes® "Ultimate" zippered encasement - zipper must be kept closed). Some insecticides may need to contact bed bugs directly to be effective. Incorporating nonchemical controls including vacuuming, low moisture steaming (AmeriVap, Hi-Tec Cleaning Systems, etc.) laundering, and removing of infested items maybe necessary to manage bed bugs and may be more important as resistant bed bug population are encountered. Reducing clutter will reduce callbacks. Fumigation can be conducted on entire buildings or certain materials, including box springs and mattresses, placed in a permanent or temporary chamber. Fumigation may be a viable option when multiunit dwellings (hotels, apartments and dormitories) have widespread infestations, when occupants are highly allergic to bites, or repeated crack and crevice applications (liquids, dust and aerosols) have failed. Fumigation does not provide residual control. Wrapping and taping mattresses in black plastic and placing in sun does NOT heat the mattress evenly and does NOT reach the bed bug's lethal threshold on the underside. See UT Extension publication PB 1763 for details. For additional bed bug information see http://eppserver.ag.utk.edu/personnel/Vail/BEDBUG%20WEB%20SITES.pdf</p>

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
BOOK LICE	Small, soft-bodied, cream-colored to grayish or light brown, wingless, fast-moving. Feed on molds, fungi. Found in books, cereals, wallpaper, boxes. May damage starched things.		Large numbers of book lice develop under excessive humid conditions, moldy books, papers, bags or cereals. Dry out infested areas. Destroy infested material of little value. Space sprays of pyrethrins may cause the book lice to scatter throughout structure.
BOXELDER BUGS See SP341-H	Flat, ½ inch long, 1/3 inch wide, dark brownish-black, with 3 lengthwise red stripes behind the head. Wings leathery at base. Membranous at tip with red veins; abdomen is red. Nymphs are smaller, wingless and bright red.	Treat listed sites on label when bugs are first seen. Tempo SC Ultra Tempo 20 WP Dragnet SFR Demand CS Demon TalstarOne Suspend SC Deltadust	These insects are attracted to buildings in the spring and fall. Large numbers collect on siding, around doors, sunny walls and attics. Use exclusion practices before pests become apparent. Inside, vacuum into a dry vac. Avoid use of space sprays if bugs found in wall voids. Dead bugs in wall voids could serve as carpet beetle food. Eliminate female (seed-pod-bearing) boxelder. Outside: vacuum the bugs into water mixed with 1 teaspoon of a liquid detergent per gallon of water in wet/dry vacuum cleaner tank.
CARPENTER BEES See SP341-P	<p>½ to 4/5 inch long with a blue-black, green or purple metallic sheen. Color and size resembles a bumble bee, but the top of the abdomen is hairless.</p> <p>These bees chew a circular, 3/8-inch entrance hole into wood and nesting gallery 4-6 inches long at a right angle to the entrance hole. Galleries used for several years may extend 10 feet.</p> <p>Carpenter bees overwinter in previously used galleries, so the structure should also be inspected in the fall. Repaint sealed area to prevent woodpecker damage.</p>	Apply dusts into the gallery openings: Tempo 1D DeltaDust Apicide <u>Sprays:</u> Tempo SC Ultra (spray or foam) Tempo 20WP (spray or foam) Cy-Kick Dragnet SFR Suspend SC 0.06% Premise .05%-0.1% (spray or foam)	Nonchemical or preventive controls include painting or varnishing wood surfaces. Use a straight wire to break up cells in tunnels. Individual bees can be caught with a net and killed. In the spring, apply foam, spray or puff of insecticidal dust into nest holes in the evening when the carpenter bees are at rest. Allow bees access to the nest for 24 hours. Seal the hole with putty, 3/8 inch diameter dowel or cork to prevent reinfestation.
CARPET BEETLES Black Carpet Beetle See SP341-I Common; or Furniture; or Varied carpet Beetles	<p>Adult 1/8 inch to 1/4 inch in length; black; brown legs. Larva 3/8 inch in length; carrot-shaped; covered with short hair and has long terminal bristles.</p> <p>Adults 1/8 inch long with white and orange; or yellow, white and black; or white, brown and yellow spots; larva with long black to brown hairs.</p>	Treat cracks, crevices and hidden area of walls, closets, stored materials, under carpets, etc. Do not apply insecticide to clothing. See precautionary statements about pesticides staining carpets. Tempo 20WP Tempo SC Ultra Cy-Kick Demand CS Dragnet SFR Suspend SC <u>Insect growth regulators:</u> Gentrol IGR Concentrate Gentrol Point Source Archer	Carpet beetles infest carpeting, clothing, fur, upholstered furniture, books, bird nests, milk powders, articles of animal products, feathers, wool, silk and other materials of animal origin. Locate food source and remove, if possible. Use good housekeeping such as cleaning floor and carpets regularly. Dry clean clothes regularly. Stored materials subject to damage should be thoroughly cleaned and stored in tight container with moth crystals.

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
CENTIPEDES	Grayish creature with long feelers and many long, slim legs. Fast moving. Long antennae.	<u>Indoors:</u> DeltaDust <u>Outside Perimeter:</u> Demon WP Cynoff WP DeltaDust Tempo 20WP Cy-Kick TalstarOne Sevin (may stain surfaces)	Feed on insects. Can bite. Usually not numerous. Active at night. Non-chemical control: leave a 12-18 inch bare zone next to foundation base. Move wood piles and other clutter away from structure. Spot-treat cracks and crevices, door thresholds and moldings where pest may crawl or as directed by label. Residuals may not provide total control. Treatments may need to come into direct contact with pest.
CHIGGERS	Very tiny mites which get on the person and cause blotches and itching. Don't invade homes, but get in yards and wooded areas.	Use commercial repellents around ankles and waist for personal protection. Apply deet repellent to skin; and Permanone 0.5% spray to shoes, cuffs and socks 2 hours before wearing. Apply sprays to infested areas of turf. Tempo SC Ultra Tempo 20 WP	Mow lawn regularly. Remove weeds and brambles from fence rows.
CLOTHES MOTHS See SP341-J	Brownish moths, wingspread 1/2" long. Larvae are 1/16-1/3" long. Gray silken cases or webbing over surface of fabric.	Vacuum prior to treatment and to remove potential food sources. Apply sprays to cracks and crevices in closets and spot treat other infested areas. Tempo 20 WP Tempo SC Ultra Cy-Kick DeltaDust Suspend SC	Adults are not attracted to light and will fly to dark corner when disturbed. Usually found on infested materials, wool, fur, feathers, hair, upholstered furniture, nonsynthetic carpets, dust and lint. Do not spray clothes. Store susceptible items with moth crystals in sealed storage container. Heed warning about staining clothing.
CLOVER MITES	Tiny (1/30 inch) mites, brown to olive green with pair of long front legs.	Apply sprays to points of entry such as foundations, windows and doors. Tempo SC Ultra (as for perimeter pests) Tempo 20 WP(as for perimeter pests) Cy-Kick CS PT 221L TalstarOne	Invade homes from the yard in great numbers in fall and spring. Stain walls or fabrics reddish-brown when crushed. Keep grass and shrubs from direct contact with house. Good weed control in turf and a vegetation free border of 12-18 inches around home will help.

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
<p>COCKROACH PB1024</p> <p>German Cockroach</p> <p>Brown Banded or Furniture Cockroach</p> <p>Oriental Cockroach</p> <p>American Cockroach</p> <p>Smokybrown Cockroach</p>	<p>About 5/8" in length, pale brown or tan with 2 parallel dark streaks on pronotum. Usually most abundant in the kitchen and bathrooms.</p> <p>1/2" to 5/8" in length, dark brown with 2 pale bands traversing wings. Widely distributed throughout the house in walls, closets, furnishing, in appliances, but abundant in kitchens.</p> <p>1 1/4" in length, dark reddish-brown to black, wings do not surpass end of abdomen. Usually found in lower floors, outdoors or in crawl space. Frequents water meters, floor drains or moist, dark areas.</p> <p>1 1/2" in length, reddish-brown with pale yellow band around pronotum. May be found throughout house, outdoors, in crawl spaces, sewers, water meters and garbage cans.</p> <p>1 to 1 1/2 inches, uniform, very dark brown to black; head shield is a solid, dark color. Takes harborage in moist, warm and dark places liketreeholes, mulches, soffits in attics with poor ventilation.</p>	<p>Do not spray repellent insecticides near baits.</p> <p><u>Baits in cracks and crevices:</u> Dupont Advion Cockroach Gel Bait Dupont Advion Cockroach Bait Arena Maxforce Roach Killer Small Bait Stations Maxforce Roach Killer Bait Gel Maxforce FC Select Roach Killer Bait Gel (for bait averse roaches) Maxforce FC Magnum Roach Killer Bait Gel Niban G Prescription Treatment Advance Cockroach Gel Bait Reservoir Prescription Treatment Avert Cockroach Gel Bait Formula 2 (Pressurized Bait Prescription Treatment Avert Dry Flowable Cockroach Bait Form. 1 Prescription Treatment Avert Cockroach Gel Bait Formula 3 Prescription Treatment Avert Cockroach Bait Stations Formula 1 Seige Gel</p> <p><u>Baits for large roaches:</u> Dupont Advion Cockroach Gel Bait Maxforce Granular Bait Maxforce Roach Killer Bait Gel Maxforce FC Magnum Roach Killer Bait Gel Niban G Seige Gel</p> <p><u>Insect Growth Regulators for small roaches</u> containing hydroprene (Gentrol, Gentrol Point Source) or pyriproxyfen (Archer, Nylar or others). IGRs are slower acting but longer lasting - sterilizes adults and prevents immatures from completing development.</p> <p><u>Lightly dust voids with:</u> PT Tri-Die Drione DeltaDust NiBor-D</p> <p><u>Sprays:</u> Tempo 20WP Tempo SC Ultra Cy-Kick Demon WP Demand CS Suspend TalstarOne</p>	<p>Prevent access to food, water and shelter. Practice good sanitation in food handling, storage and eating areas. Control moisture, prevent leaks or condensation. Seal off harborage sites such as cracks and crevices with caulk, etc. Also use exclusion practices to prevent cockroach movement.</p> <p>Use glue boards or sticky traps placed along edges in dark places to locate and monitor cockroach populations.</p> <p>Baiting is the preferred method for cockroach control. Apply baits to cracks, crevices, pipe opening into walls, joints of furniture and cabinets, pipe conduits, and elsewhere as indicated by glue board catches.</p> <p>If you chose to spray, use precautions to keep chemicals out of food, spices, and off dishes or eating utensils. Do not apply sprays where electrical shorts may occur; use baits or dusts in these areas.</p> <p>Do not use sprays when baiting because cockroaches may be repelled from the baits. Read label carefully; some products may not be labeled for food handling areas.</p>

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
CRICKETS	Black, jumping insects with long antennae. Cave or camel crickets are humped and brown.	<u>Baits:</u> Maxforce Fine Granule Insect Bait, Maxforce Granular Bait, Niban Granular Bait <u>Sprays and dusts:</u> Tempo 20WP Tempo SC Ultra Cy-Kick Demon WP Cynoff WP Demand CS TalstarOne DeltaDust Suspend SC	Black cricket may damage clothing. Repeat treatment as needed. Spray entry points into structure. Dust crawl space. Camel crickets infest damp basements, under slabs and crawl spaces. Ventilate or dry these areas. Active at night. Apply sprays into cracks and crevices where crickets dwell. Use exclusion practices. Glue boards can be used indoors around entry points and other places in basements, etc.
EARWIGS	Easily identified by pair of "pinchers" at end of abdomen.	<u>Outdoors:</u> TalstarOne Demon W Cynoff WP Dragnet SFR Demand CS	Earwigs are incidental invaders into houses. They usually dwell in leaf litter, mulch and woodpiles and are common "hitchhikers" on vegetables harvested from the garden. Moving compost piles away from the house will aid pest control. Insecticidal control is usually unnecessary for this insect. If needed, spray possible entry points and mulched area around the house.

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
<p>FLEAS PB 1596</p>	<p>Small, 1/16" long, reddish-brown, wingless insect. Body compressed laterally, legs long and adapted for jumping.</p>	<p>On pet:</p> <p>CATS, KITTENS, & PUPPIES ARE MORE SENSITIVE TO INSECTICIDES!!!! Consult a veterinarian and always read label prior to treating a pet.</p> <p>Veterinarian supplied products - usually kill fleas within 12 - 36 h or sooner and provides 90 - 95% control for about 30 days: see pesticide recommendation at http://www.ent.uga.edu/pmh/</p> <p>under <i>Homeowner</i>, and <i>Pets & Honey Bees</i>, for a thorough listing of veterinarian supplied on-pet products.</p> <p>Indoor Premise C & C Treatment Spot-treat infested areas and pet resting areas inside with following: Insect Growth Regulators [and adulticides]:</p> <p>pyriproxyfen (Archer and others)</p> <p>pyriproxyfen [and permethrin or pyrethrin] (PT Ultracide and other ready-to-use products)</p> <p>methoprene (Precor IGR Concentrate)</p> <p>methoprene [and pyrethrins or permethrin] (Precor 2000 Premise Plus Spray and other ready-to-use products)</p> <p>Adulticides Suspend SC, others</p> <p>Outdoors (when specified on label): pyriproxyfen (Archer and others) Demand CS (outdoors only) Suspend SC Talstar One Talstar PL Granular</p>	<p>Keep pets and people out of treated area (indoors and outdoors) until spray dries.</p> <p>See precautionary statements about pesticides staining carpets at the end of this bulletin.</p> <p>Step 1. With veterinarian supplied products that are currently available, control of fleas in small- to moderate-sized infestations is likely to occur by using those pet treatment products alone. May take 2 months to completely break flea life cycle.</p> <p>Sprays of pyrethrum and pyrethroids may not work as well as the newer chemistry used in pet treatments because of the potential of insecticide resistance.</p> <p>If pet treatment alone does not provide sufficient control, initiate a complete control program by April.</p> <p>Step 2. Vacuum infested areas twice a week and prior to treatment to remove eggs, larvae, adults and organic matter. Steam- cleaning carpet may also reduce populations. Eliminate fleas from pets, bedding and premises before departing on vacation.</p> <p>Step 3. Treat pet resting areas indoors and clean or remove pet bedding on the same day. Insect growth regulators important to break flea life cycle. A combination of an insect growth regulator and an adulticide may be the most efficient formulation to use.</p> <p>Step 4. Mow grass, keep weeds down and trim shrubs to expose flea eggs and larvae to lethal dessication. Irrigating areas surrounding buildings, but not against building, may kill fleas by drowning. If fleas are surviving outdoors, apply insecticide to labeled areas.</p>
<p>FLIES Face flies, cluster flies, and blue bottle flies</p>	<p>Adult flies of these three species hibernate in attics and wall voids. Cluster flies about 1/3" long, dark gray, with checkered black and silver abdomen, with gold hairs on thorax of newly emerged adults. Face fly similar in appearance to the house fly. Adult blue bottle flies have a dull gray thorax and a shiny blue abdomen.</p>	<p>Exclude flies in the fall by sealing entry points, screening behind all vents, sealing holes in walls and attics prior to pest entry in fall.</p> <p>Can apply pyrethroids to potential entry points prior to pest entry. Vacuum or use pyrethrin sprays to kill exposed flies. Use black light trap with sticky surface.</p> <p>Dust voids (boric acid not very effective) where flies may be over-wintering.</p>	<p>Cluster fly larvae are parasites of earth worms.</p> <p>Face fly larvae develop in cow patties. Adults overwinter in attics and wall voids.</p> <p>Blue bottle fly larvae develop in garbage, decaying meat, dead animals, fish and excrement.</p>

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
House flies	About 1/4" in length, dull gray color with 4 longitudinal dark stripes on the thorax.	<p><u>Treat outdoor areas where flies rest such as garbage collection sites:</u> Tempo 20 WP, Demon WP Demand CS, Dagnet SC, Suspend SC</p> <p><u>Bait:</u> Scatter, place in station or paint on. For use around commercial facilities. Maxforce Granular Fly Bait Maxforce Fly Spot Bait</p> <p>If needed, use pyrethrin space spray for temporary relief inside home.</p>	Larvae develop in warm organic matter of animal or vegetable origin. Remove trash at least twice a week to reduce fly populations in homes. Screen windows and doors. Garbage cans should have tight-fitting lids. Use insect light traps indoors. Sanitation is very important.
<p><u>Bottle Flies</u> Green Bottle Fly</p> <p>Blue Bottle Fly</p> <p>Bronze Bottle Fly</p> <p>Black Bottle Fly</p>	<p>1/2" in length; green metallic color</p> <p>1/2" in length; blue metallic color</p> <p>1/2" in length; bronze metallic color</p> <p>1/3" in length; shiny grey thorax and dull blue metallic abdomen.</p>	Treat outdoor areas where flies rest. See house fly	<p>Bottle flies indoors often indicate a dead mouse or other animal in wall voids, attic, basement, etc.</p> <p>Dispose of dead animal carcasses, animal excrement, etc.</p> <p>Garbage cans should have tight-fitting lids.</p>
Fruit Fly or Vinegar Fly	1/8" long, red eyes, tan head and thorax, abdomen gray-black.	Sanitation. Remove larval food sources. Check mops and brooms too. Use foam or other application of microbials or botanical drain cleaners/ insecticides to labeled sites (which may include drains, baseboards, behind bars, drip trays, bundles of sticky syrup lines, grout ruts, under ice machines and other equipment), that should be cleaned, but are often neglected and difficult to clean. May need to be repeated. Traps may reduce adult fly populations. Pyrethrins for temporary relief of adults.	Egg to adult in 8-11 days. Larvae in decaying fruit, vegetables and garbage cans, etc. Adults around larvae.
Moth Fly Sewer Fly	Small, scaly or hairy, long-legged moth-like flies.	Sanitation. Remove moist organic materials, clean drains with wire brush or steam clean. Following initial mechanical cleaning in commercial accounts, botanical, microbial and/or enzymatic drain cleaners can be applied to labeled sites, which may include drains, waste water traps, etc. to maintain clean surfaces. A foam formulation may work best on vertical surfaces. Pyrethrins for temporary relief of adults.	Adults rest on walls or foliage. 3-4 weeks from egg to adult. Larvae found in slimy drains, sewer backup or leaks, unsanitary garbage cans, potted plant saucers, baths or feeders for birds, clogged roof gutters or storm drains, drip lines from air conditioners, moist compost, septic tanks and other places that hold very moist organic solids.
FUNGUS GNATS See SP341-C	Adults 1/8 to 1/4 inch long. Slender larvae have shiny black head and white thread-like body.	Pyrethrins for temporary relief of adults. See SP 341-C for list of products labeled for adult and larval fungus gnats in interiorscapes. Need a category 3 certification and working under someone licensed in HRI to make interiorscape pesticide applications.	Avoid over-watering plants, let soil dry out somewhat between waterings so larvae don't have fungi to feed on. If plants not the problem, look for water leaks or other moisture problems. Check flat roof and piles of pet bird droppings which can support fungal growth. Remove moisture or dry moist areas. May enter from outdoors. Rake and reduce mulch depth to 2 - 3 inches. Adults attracted to light. Collect in windows and soil in potted plants. Run rapidly over surface.

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
HEAD LICE See SP341-S	Tiny, flat insects which infest people and clothing.	Premise sprays are not recommended for head lice control. Several louse shampoos and other hair products are available for homeowner use.	Wash infested clothing and bedding with strong soap and very hot water; tumble dry on high heat. Dry clean woolens. Do not share hair brushes, caps, etc. Use special combs to remove nits (eggs). Nits hatch by 10 days, so another application of head lice shampoo may be needed at this time. Follow label directions.
LADY BEETLE, MULTICOLOR ASIAN SP503-C	Multicolored Asian Lady Beetles (MALB) start searching for overwintering sites, your home, on the first or second day when temperatures are greater the 65°F after a dramatic drop in temperature, usually to near freezing. This usually occurs about the third week in October.	Demand CS Suspend SC TalstarOne other pyrethroids	1. Pest-Proof: seal entry points before beetles arrive. 2. Treat roof lines or soffits, vertical contrast areas, and entry points (around the following outdoor items: windows, doors, vents, pipe penetrations) with pesticides before the beetles arrive. 3. Remove dead beetles as they pile up because they may cause other MALB to aggregate. 4. If the beetles make their way into the home, vacuum or try a light trap.
MICE See PB1624	Adults weigh about ½ ounce. Dusky gray color, slender body, prominent ears, tail about as long as head and body.	Place snap traps, multiple catch traps and glue boards along paths traveled by mice. Traps or glue boards should be placed every 8-12 ft. Traps can be baited with: whole nuts, peanuts or peanut butter, dry rolled oatmeal, bacon squares, small wads of cotton or gumdrops. Baited traps should be set at right angles to rodent runs. Place trap at right angles to rodent pathway with trigger part of trap against the run.	Mice move in from outdoors in fall as temperatures decline. Exclusion practices needed, mice can fit through an opening 1/4" in diameter. Sanitation: remove access to food, water and shelter. Rodents use edges of walls, studs and pipes as guidelines. Remember to set traps where children and pets will not be hurt. Mice are curious and will normally approach traps the first night. If you don't catch a mouse in the first few nights, the trap is in the wrong location. EPA has changed allowances regarding use of rodenticide baits in the urban environment. Check for latest updates before using baits.
MILLIPEDES	Slender, brownish, multi-legged, hard-shelled, 1-2" long. Two pair of legs per body segment. Invade home from outdoors. Harmless.	<u>Outdoors:</u> Tempo 20 WP Cy-Kick Tempo SC Ultra Demon WP Cynoff WP Demand CS SuspendSC TalstarOne Sevin (may stain surfaces)	Millipedes are not insects, so insecticides not always effective. Best control obtained when pest comes in direct contact with the insecticide. Usually occasional invaders, but may invade in large numbers. Under these circumstances, non-chemical control may be more effective: remove mulch and other clutter from near the foundation, dethatch lawns and water in the morning. Prune tree limbs to dry their habitat. Use exclusion practices. Treat entry points into structure, shady areas, ivy beds, flower beds and rock walls, leaf-litter or as directed by label. Millipedes will die within 2-3 days after entering a dry structure.

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
MITES, BIRD OR RODENT	Mites occasionally found indoors because of rodent or bird nest in, on, or near structures. Some of these mites may bite people. They are small (about the size of a period), but can usually be seen with the naked eye.	<p>Dust or spray cracks and crevices around infested area.</p> <p>Tempo 1D (as for other pests) DeltaDust (as for other pests) TalstarOne (as for other pests)</p> <p>If widespread, space sprays of pyrethrins may be necessary.</p>	<p>The first step in controlling bird or rodent mites is to eliminate the host animals and remove their nesting sites. Often, the nests will be found in the attic, around the eaves and rafters or in the gutters or chimney. Gloves should be used when handling dead animals. A respirator should also be worn when removing nest materials to avoid inhaling fungal spores and other potential disease-producing organisms associated with the droppings. See chiggers for repellents.</p>

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
<p>MOSQUITOES See SP503-B</p> <p>See Entomology and Plant Pathology News web page for a list of web sites pertaining to mosquito control and West Nile virus</p> <p>http://eppserver.ag.utk.edu/mosquitoes.htm</p>	<p>Delicate insects that bite humans and animals. Breed in water. Adults stay in shrubbery, crawl spaces, etc.</p>	<p><u>Treat standing water with:</u> <i>Bacillus thuringiensis israeliensis</i> (Bti):</p> <p>Aquabac, Teknar HP-D and G, Vectobac G, CG and 12AS</p> <p>methoprene (IGR) Altosid SBG, XR-G, Pellets, Liquid, XR- briquets, briquets, Pro-G Pre-Strike Granules</p> <p>temephos Abate Pellets Abate Tire Treatment</p> <p><u>For <i>Culex</i> mosquitoes in septic conditions, treat standing water with:</u> <i>Bacillus sphaericus</i> Vectolex CG, WDG, WSP</p> <p><u>Outside Area ULV applications by ground equipment for use by trained professional personnel:</u> Anvil 10 + 10 Biomist 3 + 15 ULV Biomist 1.5 + 7.5 Biomist 30 + 30 ULV Kontrol 30 + 30 Mosquitomist One Scourge 1.5 + 4.5 (RUP) ULV Mosquito Master 412</p> <p><u>Outside residual barrier</u> (permethrin, malathion, Tempo, Suspend, Talstar and others) can be applied to vegetation on perimeter of property that is prone to rapid infestation of mosquitoes. This kills adults feeding on nectar in day and some may act as a repellent. Spray other shady damp areas where mosquitoes rest.</p> <p><u>If needed indoors</u>, use sprays containing pyrethrins in closets, stairwells, behind and beneath furniture for temporary relief.</p> <p>A more extensive list of mosquito control products for commercial and government agencies can be found at</p> <p>http://www.ent.uqa.edu/pmh/Com_Humans.pdf</p> <p>These products have not been verified to be registered in Tennessee.</p>	<p>To reduce mosquito populations, a complete mosquito control program must be followed. See UT Agricultural Extension Service Publication, The Tennessee Mosquito Control Handbook (Pesticide Applicator Licensing Manual) for more details.</p> <p>Pesticide applicators applying mosquito control pesticides on public lands and waters need to be certified in category 8 and licensed in PHMC.</p> <p>Eliminate larval sites (standing water) around structure by unclogging gutters, emptying bird baths, children's pools, pet bowls, flower pot saucers, old tires, and other containers around home twice a week. Drain or fill low areas where water collects. Easiest to control mosquitoes in immature stage because confined to water. Treat standing water with labeled insecticide. Repair screens.</p> <p>People should wear repellents when potentially exposed to mosquitoes. Apply N,N-diethyl-m-toluamide or N,N-diethyl-3-methylbenamide called DEET to skin. American Academy of Pediatrics (2003): concentration of 10% DEET appears to be as safe as products with a conc. of 30% when used according to label. Most guidelines cite that it is acceptable to use repellents containing DEET on children over 2 years of age. Children and adults can wear clothing with long pants and long sleeves while outdoors. DEET or other repellents such as permethrin can also be applied to clothing (don't use permethrin on skin), as mosquitoes may bite through thin fabric.</p> <p>On April 22, 2005 the CDC, Center for Disease Control and Prevention, added two new active ingredients, picaridin and oil of lemon eucalyptus, as suggested repellents to prevent mosquito bites.</p> <p>Mosquito netting can be used over infant carriers. More information on repellents and their safe use can be found at http://www.cdc.gov/ncidod/dvbid/westnile/qa/insect_repellent.htm</p> <p>Reduce the number of areas where adult mosquitoes can find shelter by cutting down weeds adjacent to the foundation and in yards, and mowing the lawn regularly.</p>

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
PANTRY PESTS			
Saw-Toothed Grain Beetle	Brownish black, 1/8" long, flattened with 6 saw-tooth like projections on thorax. Feeds in a wide variety of stored products, cereals, nuts, dried fruit, cookies, candy, etc.	<p>Locate food source and discard.</p> <p>Place grains, flours, nuts and other stored products in insect proof containers when they are brought home from store.</p> <p>Pheromone traps can indicate the presence of pests and may provide control without insecticides when populations are low and pests confined.</p>	To prevent infestations:
Cigarette Beetle	1/8" long, oval, reddish brown, head not visible from above, antennae saw-like. General feeder in tobacco, seasonings especially paprika, cereal, dried flowers, and a wide variety of stored foods.	<p>Vacuum cracks and crevices and wipe down pantry to remove pests and food source. Do this before insecticidal application, also.</p> <p>Do not contaminate food, dishes, shelves or utensils with pesticides. If insecticide application desired, direct spray into cracks and crevices of storage cabinet shelves when shelves are clean and all food and utensils, etc. removed.</p>	<p>1) Inspect stored products periodically,</p> <p>2) practice good sanitation,</p> <p>3) rotate stored product use so older stores are used first and none remain in storage indefinitely,</p> <p>4) have adequate ventilation to prevent moisture buildup in storage areas.</p>
Indianmeal Moth	3/8" wing span, inner 2/3 of wing grayish, outer 1/3 of wing copper in color. Feeds in coarse grain products, chocolate, nuts, dried fruit.	<p>Treat only cracks and crevices in the pantry. Do not wash off insecticide residue. Cover with paper if desired.</p>	<p>5) Insect proofing; use insect-proof package or storage procedures wherever possible.</p> <p>6) Pheromone traps can indicate the presence of pests and are available for: Indianmeal moth, saw-toothed grain beetle, confused and red flour beetle, cigarette beetle, drugstore beetle, clothes moths and others.</p>
Rice Weevil	1/8" long. Long snout on head, dark brown with 4 bright spots on wing cover. Feeds on grains.	<p>pyrethrins Tempo SC Ultra Cy-Kick DeltaDust Suspend SC Gentrol Point Source</p>	<p>Non-chemical control:</p> <p>Either destroy the infested products or salvage them by super heating to 140 degrees F for ½ hour, or super cooling in a deep freeze at 0 degree F for 4 days.</p>
Confused Flour Beetle	1/8" long, reddish-brown convex, oval shape, antennae gradually enlarged to end in a club. Cannot fly. Feeds in flour and cereal products.		Store insect-free beans in containers with tight lids.
Red Flour Beetle	1/8" long, reddish-brown convex oval shape, antennae has a distinct 3-segmented club. Feeds in flour and cereal products.		

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
<p>POWDERPOST AND OTHER WOOD-BORING BEETLES See E&PP #391</p> <p>Powderpost Beetles</p> <p>Lyctid powderpost beetle</p> <p>Anobiid powderpost beetle</p> <p>Roundheaded borers</p> <p>Old house borers</p> <p>Others</p>	<p>Shot-sized holes along with flour-like powder indicate these beetles.</p> <p>Attacks hardwoods such as oak, ash and hickory found in wood paneling, molding, window and door frames, plywood, hardwood floors and furniture. Antennae with 2-segmented club. Head protrudes forward. Reinfests seasoned wood less than 7 - 10 yrs. old.</p> <p>Attack hardwoods and softwoods. In addition to above, they also attack beams, rafters, joists, studs and other structural framing. Infestations found in moist, poorly ventilated areas such as crawl spaces, basements, etc. Head hidden by pronotum. Reinfest seasoned wood that may be decades old.</p> <p>Presence indicated by large hard-shelled beetles with long antennae.</p> <p>Broadly-oval 1/4" emergence hole made by old house borer. Larvae in tunnels packed with frass; 3 eye spots to left and right of mandibles. Beetle 3/4 inch long, grey-brown with 2 patches on wing covers; 2 bumps on thorax. Reinfests seasoned softwoods (pine).</p> <p>Neat 1/2" holes may appear in walls where beetles emerge. Don't usually reinfest seasoned softwoods (pine).</p>	<p>Products listed in this column refer to treatment for all reinfesting wood-boring beetles listed.</p> <p>Beetles that have pupated prior to insecticide application may be unaffected and may continue to emerge. Insecticide applications should prevent reinfestation.</p> <p>Products containing disodium octaborate tetrahydrate (DOT) and glycols (Boracare, Shell-Guard), may penetrate wood further than other residuals, but penetration is variable and depends on moisture content of the wood and other factors. Other DOT products include Timbor, Armor-Guard and others. DOT treatments must be made to unfinished surfaces as they will not penetrate paint or varnish; sand or power wash logs prior to treatment. Paint, spray, inject or brush on.</p> <p><u>Other residual insecticides</u> that are injected or sprayed or brushed on to coat the surface of the wood. Spray or brush on until wet, but NOT to runoff: TalstarOne Tempo SC Ultra PT 221L (injection) Demon TC Suspend SC Dragnet SFR</p> <p>If infestation spread into walls or between floors, some of the above products can be foamed. If foaming is not feasible because of electrical sources, fumigation may be needed.</p> <p>Fumigation is costly and should only be considered as a last resort. If only small articles infested such as furniture, antiques, etc., they can be fumigated at a lower cost. Only professional pest control operators licensed to fumigate can perform this operation.</p>	<p>Determine extent of infestation. Signs for powder post beetles are: flour-like "frass" dropping from pinhead-sized or slight larger holes, Anobiids' frass are more gritty than Lyctids; adult beetles attracted to light may be found on window sills or foundation vents. Important to determine if infestation active or not. Mark or seal existing holes, vacuum existing sawdust, recheck wood for new holes in spring or early summer. These beetles damage wood slowly. If "frass" is yellow, caked or covered with dust or debris, that damage is old. Old house borers can be detected by hollow sound when wood tapped.</p> <p>Prevention:</p> <p>1) don't use old lumber from a barn or wood pile unless it has been treated</p> <p>(2) don't use improperly dried or stored lumber</p> <p>(3) inspect firewood prior to bringing into structure</p> <p>(4) paint, varnish or otherwise seal wood to prevent exposed edges</p> <p>(5) seal previous emergence holes to prevent egg-laying sites.</p> <p>New houses usually infested by use of infested lumber. May also come from firewood.</p> <p>Alternative controls for powder post beetles: small items, such as picture frames can be heated at 120 to 140 F for six hours to kill existing life stages. Freezing (0 F) infested wood for 72 hours will also kill all life stages .</p> <p>If all evidence indicates the infestation is localized, wood could be replaced. Watch for new holes in adjacent areas.</p> <p>Decrease moisture in wood through ventilation and moisture barriers. Central heat and air may reduce wood moisture so there is insufficient moisture to support large infestations in living areas. Wood kept below 14 percent moisture would be unsuitable to Anobiid powderpost beetle reinfestation or development. Professionals have moisture meters.</p>

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
<p>RATS PB1624</p> <p>EPA has changed allowances regarding use of rodenticide baits in the urban environment. Check for latest updates before using baits.</p>	<p>Norway rat: 12-18 inches, tail shorter than head and body, body heavy and thick, ears small</p> <p>Roof rat: 12-17 inches, tail longer than head and body, body light and slender, ears larger.</p> <p>Young rat : 6-7 inches feet large, head large</p> <p>House mouse: 6-7 inches feet small, head small</p> <p>Droppings:</p> <p>Roof rat: pointed, about ½ inch</p> <p>Norway rat: blunt, about 3/4 inch</p> <p>House mouse: pointed, about 1/8 inch</p>	<p>When rats are plentiful or where unsanitary conditions exist with shelter, poisoned baits are the best control method. Often community- wide control needed. Poison baits are available as ready to use, premixed baits. They come in many forms: parafinized blocks for outdoor use and high humidity areas; treated meal; seeds; or parafinized pellets in bulk or in "place packs" for indoor use. Water baits are sold as packets of concentrate that are mixed with water. They are administered with a chick waterer and are useful in areas where rodent food is abundant. Poison baits should be placed where they are inaccessible to children and pets. Where rodent runs are exposed and in most outdoor situations, tamper proof bait boxes should be used. Vitamin K is the antidote for anti-coagulants.</p>	<p>Exclusion practices needed. Rats can fit through an opening ½ inch in diameter. Locate entrance into structure and exclude. Use materials such as galvanized, stainless or other non-rusting metal such as 24-gauge sheet metal or 19-gauge hardware cloth with 1/4 inch or smaller opening; brick, concrete block, tile or glass; steel wool with expandible foam; and others. Remove debris such as piles of waste lumber or trash, used feed sacks, abandoned large appliances and wood piles from next to structure. Store pet foods and seed in rodent proof glass or metal containers. Place snap traps, multiple catch traps and glue boards along paths traveled by rats. Of the snap traps, the expanded trigger trap is the most versatile since it can be baited. Rats are bait shy. Leave baits in place for at least a week before moving. Place trap 90 degrees to rodent pathway with trigger part of trap against the run. Rodents use edges of walls, studs and pipes as guidelines. Snap traps can be baited with: whole nuts; raisins or grapes for roof rats; sardines packed in oil for Norway rats; peanuts or peanut butter; bacon squares; or small wads of cotton.</p> <p>Often area-wide effort needed.</p>
<p>SILVERFISH AND FIREBRATS See SP341-O</p>	<p>Grayish, wingless, rapid-moving insects with 3 long tails. Feed on starchy materials such as bookbinding, wallpaper, cardboard, etc.</p>	<p>Niban Fine Granular Bait Tempo 20 WP Cy-Kick Demand CS PT565 XLO Suspend SC DeltaDust TalstarOne</p>	<p>Treat crack and crevice where silverfish and firebrats may dwell. Attics many times source of infestation.</p>
<p>SKUNKS PB1624</p>	<p>These animals many times live in the ground around or under homes.</p>	<p>Bac-Azap biological odor control or others can be applied to eliminate odors.</p>	<p>Trap and remove skunks from property. Seal the foundation to prevent entry under building.</p>
<p>SNAILS AND SLUGS</p>	<p>Long, grayish, shiny, soft-bodied creatures. Will attack various plants. Leave slime trails on walks and walls.</p>	<p>Snail and slug killer baits containing metaldehyde.</p>	<p>Remove boards and plastic or plant debris and dry damp areas adjacent to foundation.</p>
<p>SNAKES PB1624</p>	<p>Snakes of various kinds, den around or invade homes and other buildings.</p>	<p>Place a pile of cool, damp rags in building where snake was last seen. Snake will be attracted and can be removed. Large glue boards can trap snakes.</p>	<p>Mouse-proof building. Mow lawns and field to control grass, weeds and brush. Remove boards, flat rocks, trash piles and other debris.</p>
<p>SOWBUGS OR PILLBUGS</p>	<p>Grayish, hard-shelled, many-legged creatures appear on walks and patios. Roll up in ball when disturbed. Occasional invaders.</p>	<p>Chemical control usually not necessary for this pest. If needed, apply to infested areas outdoors around perimeter of structure. This stops any invasion into the house.</p> <p>TalstarOne DeltaDust Suspend SC</p>	<p>Remove leaf piles, grass clippings, old boards, wood piles and other debris from around foundation. Leave a 12-18 inch plant /mulch free zone next to foundation base. Use exclusion practices: caulk cracks around foundation and screen vents in foundation. Drain and dry area around house.</p>

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
<p>SPIDERS or SCORPIONS</p> <p>See PB 1193</p> <p>See PB 1191</p>	<p>Many kinds invade homes, basements and roof overhangs from outdoors.</p> <p>Two species most dangerous in Tennessee:</p> <p>Black Widow: dark black spider with red hour glass shape on bottom of abdomen. More of an outdoor pest along perimeter of buildings. Use outside perimeter treatment with residuals.</p> <p>Brown Recluse, light brown spider, with legs reaching to the size of a quarter or half dollar, dark violin shape on back of front portion of head, 3 pairs of eyes arranged in a semi-circle. Brown recluse pose a serious threat. Quite often a professional pest control company should be used. Remove unnecessary clutter and webs from indoors and outdoors and vacuum especially, under furniture.</p>	<p><u>Dusts:</u></p> <p>DeltaDust EcoPCO® D•X Dust Insecticide Tempo 1D</p> <p><u>Sprays:</u> Demand CS Tempo 20 WP Tempo SC Ultra Cy-Kick Demon WP Suspend SC TalstarOne</p> <p><u>Space or aerosol:</u> pyrethrins</p> <p>Brown Recluse: Use residual sprays (Demand, Tempo, Cy-Kick, Demon or others) around exterior foundation, eaves, closets, storage areas and rugs. Dusts can be applied to wall voids, attics and inaccessible crawl spaces. ULV or aerosol treatment with pyrethrins to kill exposed spiders and flush others onto surfaces treated with residuals.</p>	<p>Beneficial organisms because they feed on pest insects. Occasional invaders that can be vacuumed or swept out the door. Remove wood or mulch piles away from house to lower abundance of their insect food source.</p> <p>Apply insecticides to crawl spaces, basements, attic, eaves and outdoor areas of home. Clean up debris where scorpions and spiders hide. Replace outdoor lights with yellow bug lights.</p> <p>Scorpions will fluoresce under a black light, so they and their breeding areas can easily be seen at night. Use glue boards to trap spiders and locate infested areas. Glue boards should be placed against walls and other guidelines where spiders are suspected.</p> <p>Efforts to control brown recluse will cause spiders to become more active. Prevent bites by checking shoes and clothing before wearing, by pulling beds away from walls, and preventing bed skirting and bedspreads from touching the floor. Place glue boards under bed posts.</p>
<p>SPRINGTAILS</p>	<p>Small, jumping insects with a forked spring mechanism. A dark species, called snow fleas, may be present in the winter.</p>	<p>If pest-proofing unsuccessful, try a perimeter treatment as for other occasional invaders.</p> <p>Cy-Kick CS Suspend SC TalstarOne Sevin (may stain surfaces)</p>	<p>Usually found in moist decaying vegetation and are incidental invaders into houses. Prevent entry into home by pest-proofing. There are a few reports of large populations entering homes. Dry out surrounding landscape, water only in morning, etc.</p>

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
<p>TERMITES See PB 1344</p>	<p>Termites invade and eat wood and other cellulose material, causing extensive damage in structural parts of a building. Their presence may not be discovered until they swarm, years after infesting a structure.</p> <p>Workers and soldiers: soft-bodied insects 1/8 to 3/16 inch long.</p> <p>Swarms are black to brown with pearly wings. Swarms are easily distinguished from winged ants by termites straight antennae, broadly attached thorax to waist and four nearly equal wings.</p> <p>Inspect for signs of termite infestation: irregular earthen tubes constructed across walls, floors and foundation.</p> <p>Hammer or probe timbers with a sharp instrument. Damaged wood will be soft, channeled, unsound and may possibly reveal the termite infestation itself.</p> <p>Use a moisture meter. Active termites will increase moisture reading relative to uninfested areas.</p> <p>Termites commonly enter homes around doors, wooden steps and porches and unexcavated portions of structures. The easiest access points are where wood is in direct contact with the soil.</p> <p>(Description continued on next page)</p>	<p>Listed below are products available to professionals (Manufacturer): + F = may also be foamed</p> <p>Soil treatment: Do not apply near (within 100 ft.) any body of water, cistern, or well.</p> <p>Nonrepellents (Newer a.i.s) Termites do not detect these insecticides and hence walk over the treated soil. Termiticide may be transferred back to colony.</p> <p>chlorfenapyr Phantom 0.125, 0.25% (BASF) + F fipronil Termidor 80WG, 0.06%, 0.09%, 0.125% (BASF)+F Termidor SC, 0.06%, 0.09%, 0.125% (BASF) +F imidacloprid Premise 75, 0.05, 0.1% (Bayer)+F Premise 2 (small jobs and foaming) (Bayer) Premise 0.5 SC (in-line injector) 0.05, 0.1% (Bayer)</p> <p>Pyrethroids (Older a.i.) In general, this groups tends to be repellent, thus treatments must be applied to create a continuous barrier. Many generic pyrethroids are now on the market.</p> <p>bifenthrin TalstarOne 0.06, 0.12% (FMC) + F</p> <p>cypermethrin Demon TC 0.25, 0.5, 1% (Syngenta)+F Prevail FT 0.25, 0.5, 1% (FMC) +F</p> <p>permethrin Dragnet SFR 0.5, 1, 2% (FMC) +F Prelude 0.5, 1, 2% (Syngenta) +F</p> <p>Combination product Transport 0.11% bifenthrin and acetamiprid (FMC)</p> <p>A more complete list of termiticides is available at http://www.flaes.org/pdf/TermiticidesRegisteredInFlorida.pdf , but not all FL termiticides listed may be registered in Tennessee.</p> <p>(Wood treatment continued on next page)</p>	<p>Follow correct construction practices.</p> <p>This is the best protection against a termite infestation:</p> <ol style="list-style-type: none"> (1) Remove all wood materials from around and under the house; (2) Remove all form boards and construction stakes; (3) Construct a termite-proof foundation; (4) Have at least 30 in. of clearance under buildings; (5) Have proper ventilation and light under all parts of the building; (6) Use a moisture barrier in crawl spaces; (7) Drain water away from building; (8) Have no wood in contact with the ground, or treat those timbers that require ground contact with approved preservatives/borates (9) Make periodic inspection of buildings. <p>Find a reputable professional to treat.</p> <p>Collect some swarms for identification and vacuum rest. Leave mud tubes in place until professional pest control applicator arrives.</p> <p>Newer Exterior Perimeter/Localized Interior treatment (EP/LIT) supplement to Termidor Labels. Ensure localized interior treatment is made. Label and more information at www.pestcontrolfacts.com . Premise also has an Exterior Perimeter/Interior Spot Treatment, see http://www.backedbybayer.com/bayer/crops/cience/backedbybayer.nsf/id/EN_Pest_Labels_MSDS for specifics.</p> <p>Tennessee Department of Agriculture now prefers the use of a disclosure form if less than a complete treatment is applied. Applications following the Exterior Perimeter/Interior treatments of Termidor or Premise are now considered a full treatment.</p> <p>New NPMA 33 Wood-destroying Insect Infestation report required for most real estate transactions. Sample form and directions are found at www.pestworld.org</p>

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
<p>TERMITES Cont'd</p>	<p>Some termiticide labels have listed variable rates depending on soil type.</p> <p>To ensure your home is treated as directed by the label, ask the professional to provide you with a copy of the label.</p> <p>Suggested volumes listed below are usually for the lowest rate.</p> <p>In general, horizontal barriers (under slab) should receive 1 gallon of diluted termiticide per 10 square feet or 1.5 gallons if coarse fill.</p> <p>Vertical barriers (along both sides of foundation wall, around plumbing, piers and conduits) should receive 4 gallons of dilution per 10 linear feet per foot of depth (into a trench 6 inches wide and 1 foot deep or) to the top of the footing.</p> <p>Voids in hollow masonry foundation walls should be treated at a rate of 2 gallons per 10 linear foot so the dilution will reach the top of the footing.</p> <p>Occasionally, moisture damaged wood in roofs can support an aerial infestation. No mud tubes will reach to ground. Attic inspection is important, too.</p>	<p>Wood treatment: treat galleries and wall voids; spray or brush on until wet, but NOT to runoff. Used to supplement a soil treatment. Most termiticides can be used in this manner.</p> <p>disodium octaborate tetrahydrate can be applied to wood as a pretreatment barrier or as a second barrier (see labels for sites and more details). Paint, spray, foam, inject or brush.</p> <p>Bora-Care (extensive label, other sites in addition to wood can be treated)</p> <p>Jecta (injection, for wood in contact with the ground) and others.</p> <p>Baits: Termites feed on bait and spread bait to colony to eliminate or suppress it. See PB1344 for advantages and disadvantages of baits.</p> <p>Sentricon® Colony Elimination System (DowAgrosciences LLC): Bait (Recruit IV and Recruit IV AG) contains a chitin synthesis inhibitor, noviflumuron..</p> <p>Hex-Pro™ Temite Baiting System (DowAgrosciences LLC): Bait (Shatter) contains a chitin synthesis inhibitor, hexaflumuron.</p> <p>Exterra Termite Interception and Baiting System (Ensysstex, Inc.) contains a chitin synthesis inhibitor. diflubenzuron .</p> <p>Advance™ Termite Baiting System (Whitmire-MicroGen) contains a chitin synthesis inhibitor. diflubenzuron .</p>	<p>Effective control measures for a soil treatment should include:</p> <ol style="list-style-type: none"> 1) Inspect basement and underside of house thoroughly to determine the area and extent of infestation. 2) Inspect attic for termite tubes and damage to joists, rafters, flooring and stored materials. 3) Disrupt and block all termite tubes (unless baiting) 4) Ditch the entire foundation inside and out and treat the soil replaced in the trenches with chemicals. 5) Repair all foundation and basement floor and wall breaks with rich concrete. 6) Break all wood-soil contacts, treat such areas with chemicals. 7) Treat infested timbers and replace those which are badly infested. 8) Treat hollow spaces in the foundation - concrete blocks, piers, chimney bases, spaces behind brick veneer. Ditch and treat inside of porch foundations, under patios, under concrete slabs and the surface of ground under porches and similar dead places. 9) Provide ventilation and drainage beneath house and porches. 10) Remove all scrap wood from beneath house.

PESTS	DESCRIPTION	CONTROL MATERIALS AND METHODS	REMARKS
<p>TICKS See PB 726</p>	<p>Grayish or brown, round, hard-shelled, 8-legged creatures which invade homes, yards and get on pets and people.</p>	<p>Insecticide applications are most effective when directed into areas where ticks and their animal hosts are likely to frequent. Pay particular attention to borders and fences between wooded or brushy areas and the lawn, around ornamental plantings, beside foot paths, house and dog house. Allow surface to dry before people or pets have access.</p> <p>Tempo SC Ultra Tempo 20 WP Talstar EZ granular Talstar F TalstarOne and other synthetic pyrethroid insecticides.</p> <p><u>Indoors for brown dog tick:</u> Cynoff WP Demand CS TalstarOne Cy-Kick other pyrethroids</p> <p><u>Repellents:</u> Apply deet to skin; and Permanone 0.5% spray to shoes, cuffs and socks 2 hours before wearing.</p> <p><u>Dogs:</u> fipronil (Front Line) spot-ons available from veterinarians</p> <p>BioSpot and other permethrin containing spot-ons</p> <p>collars containing amitraz (don't use around small children or dogs that may chew collar)</p>	<p>Nonchemical methods for reducing tick problems include mowing the lawn and controlling weeds.</p> <p>This has three advantages - it lowers the moisture in the grass microclimate and allows sunlight to penetrate, which tends to cause ticks to dry out; it discourages rodents (which may serve as hosts) from nesting; and lastly, because there is less plant matter, less pesticide may be needed if a treatment is necessary. Also, removing debris, weeds or clutter from around the house discourages rodents from nesting.</p> <p>Repair entry points into the house to discourage possible tick hosts from entering. Cracks and crevices, both indoors and out, can be sealed to reduce hiding places for ticks. Inspect and clean pets and their bedding frequently. If bedding is infested, it can be cleaned or destroyed.</p> <p>In the home, ticks stay around baseboards and walls.</p> <p>Use insecticides in cracks and crevice in the home for brown dog tick.</p>
<p>WASPS, HORNETS, YELLOW-JACKETS See SP290-A, SP341-M</p>	<p>Many types build paper and mud nests around homes, in ground or in shrubs.</p>	<p><u>Dusts:</u> Tempo 1D DeltaDust Drione Apicide</p> <p><u>Sprays:</u> Bee and wasp killer aerosols Tempo 20WP Tempo SC Ultra Demon WP Eco PCO Jet</p> <p>Victor Yellow Jacket Trap</p>	<p>Wait until dark when wasps return to nest and are slow due to cooler temperatures. Apply insecticides to nest opening and seal nest opening if possible.</p> <p>Remove mud nests in winter to destroy overwintering forms.</p> <p>Traps can used to reduce foraging yellow jacket populations. Place away from areas people congregate.</p>

For Pest Management Professionals

Trade Name	Chemical Name or Use	Website
Abate Tire Treatment, Pellets	temephos	Clarke Mosquito Control http://www.clarkemosquito.com
Advance Termite Bait System	diflubenzuron	Whitmire Mirco-Gen http://www.wmmg.com
Altosid Briquets (B), Liquid Larvicide, Extended Residual Briquets (XRB), Pellets (P), Extended Residual Granular (XRG), Single Brood Granular (SBG), Pro -G	methoprene	Wellmark http://www.altosid.com/
Anvil 10 + 10	sumithrin and PBO	Clarke Mosquito Control http://www.clarkemosquito.com
Apicide	carbaryl	Mystic Chemical Company http://www.apicide.com/Apicide_pail_label_may_08.pdf
Archer	pyriproxyfen	Syngenta http://www.syngentaprofessionalproducts.com/ppm
Armor-Guard	disodium octaborate tetrahydrate	NovaGuard http://www.novaquard.com
Bac-Azap	enzyme-producing bacteria and others	Nisus http://www.nisuscop.com/
Bedlam	sumithrin® MGK 264	McLaughlin, Gormley, King Co. http://www.mgkpro.com
Biomist 3 + 15 ULV, 1.5 + 7.5 ULV, 30 + 30 ULV	permethrin and PBO	Clarke Mosquito Control http://www.clarkemosquito.com
Bora-Care	disodium octaborate tetrahydrate	Nisus http://www.nisuscop.com/
Cy-Kick	cyfluthrin	Whitmire-Microgen http://www.wmmg.com/products/index_mkt_spc.asp
Cynoff 40 WP	cypermethrin	FMC http://pestsolutions.fmc.com/
Cynoff EC	cypermethrin	FMC http://pestsolutions.fmc.com/
CB-80	pyrethrin, PBO	CB Professional Products http://www.cbproproducts.com
D-Force HPX	deltamethrin	CB Professional Products http://www.cbproproducts.com
DeltaDust	deltamethrin	Bayer http://www.backedbybayer.com/bayer/cropscience/backedbybayer.nsf/id/EN_Pest_Labels_MSDS
Demand CS	lambda cyhalothrin	Syngenta http://www.syngentaprofessionalproducts.com/ppm
Demon WP	cypermethrin	Syngenta http://www.syngentaprofessionalproducts.com/ppm
Demon TC	cypermethrin	Syngenta http://www.syngentaprofessionalproducts.com/ppm
Dragnet SFR	permethrin	FMC http://pestsolutions.fmc.com/
Drione	silica gel and pyrethrins	Bayer http://www.backedbybayer.com/bayer/cropscience/backedbybayer.nsf/id/EN_Pest_Labels_MSDS

Trade Name	Chemical Name or Use	Website
DuPont Advion Ant Bait Arena	indoxacarb	DuPont http://www2.dupont.com/Professional_Products/en_US/Label_MSDS_Info/index.html
DuPont Advion Ant Gel	indoxacarb	DuPont http://www2.dupont.com/Professional_Products/en_US/assets/downloads/pdfs/H65465.pdf
DuPont Advion Cockroach Bait	indoxacarb	DuPont http://www2.dupont.com/Professional_Products/en_US/Label_MSDS_Info/index.html
DuPont Advion Mole Cricket Bait	indoxacarb	DuPont http://www2.dupont.com/Professional_Products/en_US/Label_MSDS_Info/index.html
EcoPCO® D•X Dust Insecticide	2-phenethyl propionate & pyrethrins	Eco SMART http://www.ecosmart.com/Commercial/Products/main.asp
EcoPCO Jet	eugenol, 2-phenethyl propionate	Eco SMART http://www.ecosmart.com/Commercial/Products/main.asp
Exterra Termite Interception & Baiting System	diflubenzuron	Ensystem http://www.ensystem.com/
Gentrol	hydroprene	Wellmark- Zoecon http://www.zoecon.com/products.htm
Gentrol Point Source	hydroprene	Wellmark- Zoecon http://www.zoecon.com/products.htm
Gourmet Ant Bait Gel Gourmet Ant Bait Liquid	disodium octaborate tetrahydrate	Innovative Pest Control Products http://www.antcafe.com/index.html
Jecta	10% disodium octaborate tetrahydrate	Nisus http://www.nisuscop.com/
Kicker	pyrethrin and PBO	Bayer http://www.backedbybayer.com/bayer/cropscience/backedbybayer.nsf/id/EN_Pest_Labels_MSDS
Kontrol 30-30	permethrin and PBO	Univar http://www.masterline.com/pdfs/Kontrol30-30Label122105.pdf
Mattress Safe	bed encasement	http://www.mattresssafe.com/
Maxforce Ant Killer Bait Stations	1% hydramethylnon	Bayer http://www.bayerprocentral.com/pest/labels/index.html
Maxforce Ant Killer Bait Gel	fipronil	Bayer http://www.bayerprocentral.com/pest/labels/index.html
Maxforce Roach Killer Small Bait Stations	hydramethylnon	Bayer http://www.bayerprocentral.com/pest/labels/index.html
Maxforce Carpenter Ant Bait Gel	0.001% fipronil	Bayer http://www.bayerprocentral.com/pest/labels/index.html
Maxforce FC Magnum Roach Killer Bait Gel	0.05% fipronil	Bayer http://www.backedbybayer.com/Bayer/CropScience/BackedByBayer.nsf/id/EN_Pest_Maxforce_FC_Magnum
Maxforce FC Roach Bait Stations	fipronil	Bayer http://www.bayerprocentral.com/pest/labels/index.html
Maxforce FC Select Roach Killer Bait Gel	fipronil	Bayer http://www.bayerprocentral.com/pest/labels/index.html

Trade Name	Chemical Name or Use	Website
Maxforce Roach Killer Bait Gel	hydramethylnon	Bayer http://www.bayerprocentral.com/BAYER/CropScience/BackedByBayer.nsf/id/EN_Pest_Labels_MSDS
Maxforce Fine Granule Insect Bait	hydramethylnon	Bayer http://www.bayerprocentral.com/BAYER/CropScience/BackedByBayer.nsf/id/EN_Pest_Labels_MSDS
Maxforce Fly Spot Bait	imidacloprid	Bayer http://www.bayerprocentral.com/BAYER/CropScience/BackedByBayer.nsf/id/EN_Pest_Labels_MSDS
Maxforce Granular Bait	hydramethylnon	Bayer http://www.bayerprocentral.com/BAYER/CropScience/BackedByBayer.nsf/id/EN_Pest_Labels_MSDS
Maxforce Granular Fly Bait	imidacloprid	Bayer http://www.bayerprocentral.com/BAYER/CropScience/BackedByBayer.nsf/id/EN_Pest_Labels_MSDS
Mosquitomist One	chlorpyrifos	Clarke Mosquito Control http://www.clarkemosquito.com
Mother Earth Granular Scatter Bait	boric acid	Whitmire Micro-Gen http://www.wmmg.com/pdf/label/MEGSB_SPEC_v11_WEB.pdf
Niban FG = Niban Fine Granular Bait	orthoboric acid	Nisus http://www.nisuscorp.com/
Niban G = Niban Granular Bait	orthoboric acid	Nisus http://www.nisuscorp.com/
NiBor-D	disodium octaborate tetrahydrate	Nisus http://www.nisuscorp.com/
Optigard Ant Gel Bait	thiamethoxam	Syngenta http://www.syngentaprofessionalproducts.com/ppm/
Pharorid - discontinued	methoprene	
Phantom	chlorfenapyr	BASF http://www.pestcontrolfacts.com/
Precor Plus 2000 Premise Spray	permethrin, methoprene, phenothrin, etc.	Wellmark - Zoecon http://www.zoecon.com/products.htm
Precor IGR Concentrate	methoprene	Wellmark - Zoecon http://www.zoecon.com/products.htm
Premise 75, 2, and 0.5	imidacloprid	Bayer http://www.bayerprocentral.com/
Prelude	permethrin	Syngenta http://www.syngentaprofessionalproducts.com/ppm/
Prescription Treatment Advance 375A Select Granular Ant Bait	0.011% abamectin	Whitmire Micro-Gen http://www.wmmg.com
Prescription Treatment 388B Advance Ant Bait Gel	5.4% sodium tetraborate decahydrate (borax)	Whitmire Micro-Gen http://www.wmmg.com
Prescription Treatment 381B Advance Liquid Ant Bait	1.3% sodium tetraborate decahydrate (borax)	Whitmire Micro-Gen http://www.wmmg.com
Prescription Treatment Advance Granular Ant Bait Formula 1	abamectin	Whitmire Micro-Gen http://www.wmmg.com
Prescription Treatment Advance Cockroach Gel Bait Reservoir	0.5% dinotefuran	Whitmire Micro-Gen http://www.wmmg.com
Prescription Treatment Avert Cockroach Gel Bait Formula 3	abamectin	Whitmire Micro-Gen http://www.wmmg.com/avert/shell.asp?id=2

Trade Name	Chemical Name or Use	Website
Prescription Treatment Avert Dry Flowable Cockroach Bait Formula 1	abamectin	Whitmire Micro-Gen http://www.wmmg.com
Prescription Treatment Avert Cockroach Bait Stations Formula 1	abamectin	Whitmire Micro-Gen http://www.wmmg.com
Prescription Treatment Avert Cockroach Gel Bait Formula 2 (Pressurized Bait)	abamectin	Whitmire Micro-Gen http://www.wmmg.com
Prescription Treatment Cy-Kick CS Controlled Release and Pressurized	cyfluthrin	Whitmire Micro-Gen http://www.wmmg.com
Prescription Treatment Perma-dust 35.5 %	boric acid	Whitmire Micro-Gen http://www.wmmg.com
Prescription Treatment Tri-Die Pressurized	silica + pyrethrin	Whitmire Micro-Gen http://www.wmmg.com
Prescription Treatment Ultracide	nylar (pyriproxyfen), pyrethrins, permethrin, etc.	Whitmire Micro-Gen http://www.wmmg.com
Prescription Treatment 221L	lambda-cyhalothrin	Whitmire Micro-Gen http://www.wmmg.com
PT565 XLO	pyrethrin, d-trans allethrin + others	Whitmire Micro-Gen http://www.wmmg.com
Prevail	cypermethrin	FMC http://pestsolutions.fmc.com/
Pro-Joe-S Ant Bait Gel Formula 4.5	4.5% boric acid	Blue Diamond Company
Protect-A-Bed with BugLock™ 3 sided zipper system and ALLERZIP™ seal (bed encasement for bed bugs)	bed encasement	Protect-A-Bed http://www.protect-a-bed.com/
Recruit IV and Recruit IV AG	noviflumuron	DowAgroSciences http://www.dowagro.com/ppm/index.htm
Scourge - Restricted Use Pesticide	resmethrin + PBO	Bayer http://www.bayerprocentral.com/
Siege gel	hydramethylnon	CB Professional Products http://www.cbproproducts.com
Sentricon Colony Elimination System	directions for use	DowAgroSciences http://www.cdms.net/manuf/1prod.asp?pd=7444&lc=0
Shatter	hexaflumuron	DowAgroSciences http://www.dowagro.com/hexpro/
Steri-Fab	d-phenothrin isoproponal & others	Noble Pine Products Company http://www.sterifab.com/home.html
Suspend SC	deltamethrin	Bayer http://www.bayerprocentral.com/
Talstar EZ granules	bifenthrin	FMC http://pestsolutions.fmc.com/
Talstar F	bifenthrin	FMC http://pestsolutions.fmc.com/
Talstar One, P	bifenthrin	FMC http://pestsolutions.fmc.com/
Teknar HP-D, G	Bacillus thuringiensis subspecies israelensis toxin	Valent BioScience http://www.valentbiosciences.com/agricultural_products/agricultural_products_5.asp
Tempo 20WP	cyfluthrin	Bayer http://www.bayerprocentral.com

Trade Name	Chemical Name or Use	Website
Tempo SC Ultra	β -cyfluthrin	Bayer http://www.bayerprocentral.com/
Termidor SC, WG	fipronil	BASF http://www.pestcontrolfacts.org/
Terro -PCO Liquid Ant Bait	5.4% borax or sodium tetraborate decahydrate	Nisus http://www.nisuscop.com/
Tim-bor	disodium octaborate tetrahydrate	Nisus http://www.nisuscop.com/
Transport GHP Insecticide	bifenthrin, acetamiprid	FMC http://www.fmcprosolutions.com/tabid/2362/Default.aspx
Transport Termiticide Insecticide	bifenthrin, acetamiprid	FMC http://www.fmcprosolutions.com/tabid/2362/Default.aspx
ULV Mosquito Master 412	chlorpyrifos, permethrin	Clarke Mosquito Control http://www.clarkemosquito.com
Vikane	sulfuryl fluoride	DowAgroSciences http://www.dowagro.com/ppm/vikane/bedbugs/options.htm
VectoBac CG, G, 12AS	Bacillus thuringiensis subspecies israelensis toxin	Valent BioScience http://www.valentbiosciences.com/agricultural_products/agricultural_products_9.asp
VectoLex CG, WSP, WDG	Bacillus sphaericus	Valent BioScience http://www.valentbiosciences.com/agricultural_products/agricultural_products_9.asp

PRECAUTIONARY STATEMENT

To protect people and the environment, pesticides should be used safely. This is everyone's responsibility, especially the user. Read and follow label directions carefully before you buy, mix, apply, store, or dispose of a pesticide. According to laws regulating pesticides, they must be used only as directed by the label.

DISCLAIMER STATEMENT

This publication contains pesticide recommendations that are subject to change at any time. The recommendations in this publication are provided only as a guide. It is always the pesticide applicator's responsibility, by law, to read and follow all current label directions for the specific pesticide being used. The label takes precedence over the recommendations found in this publication. Use of trade or brand names in this publication is for clarity and information; it does not imply approval of the product to the exclusion of others which may be of similar, suitable composition, nor does it guarantee or warrant the standard of the product. The author(s), The University of Tennessee Institute of Agriculture and the University of Tennessee Extension assume no liability resulting from the use of these recommendations.